












**MEN'S STI's (PAGE 1)**

<b>STI</b>	<b>IMAGE</b>	<b>KNOWN AS</b>	<b>COURSE</b>	<b>SYMPTOMS</b>	<b>TREATMENT</b>	<b>WITHOUT TREATMENT</b>
<p><b>Syphilis</b></p> <p>Image Source: University of Washington Seattle STD/HIV Prevention Training Centre</p>		The Pox	<p>Stage 1: 1-3 months</p> <p>Stage 2: 3-6 months</p> <p>Stage 3: Many years</p>	<p>Stage 1: A painless sore called a chancre.</p> <p>Stage 2: Fever, headache and general rash.</p> <p>Stage 3: Very ill. The cause is not easy to find.</p>	Antibiotics	Severe infection, infertility, skin diseases, arthritis
<p><b>Gonorrhoea</b></p> <p>Image Source: University of Washington Seattle STD/HIV Prevention Training Centre</p>		Drip, Clap, Drop	1 - 10 days	Painful urination, discharge or drip from penis.	Antibiotics	Severe infection of penis and reproductive organs
<p><b>Trichomoniasis</b></p>		Trich	1 week	May experience thin and whitish discharge from the penis; pain/burning sensation when passing urine; inflammation of the glands and foreskin (redness)	Antibiotics	Fever, infection of the genital organs.
<p><b>Chlamydia</b></p> <p>Image Source: Southern Mississippi University Health.Promotion@usm.edu</p>		PID	1-3 weeks	Discharge from penis, painful urination	Antibiotics	Severe infection of reproductive organs, infertility
<p><b>Non-Gonococcal urethritis</b></p>		NGU	1-3 weeks	Painful and frequent drip or discharge from the penis it's an infection of the prostatitis, epididymis or urethral stricture	Antibiotics	Infertility
<p><b>Pubic Lice</b></p> <p>Image Source: University of Washington Seattle STD/HIV Prevention Training Centre</p>		Crabs	Immediately	Itching in area of the chest, genital hair. Lice crawling and small eggs on hair or clothing.	Special shampoos or lotions. All clothing and bedding must be washed in hot, soapy water.	Irritation, secondary infection due to scratching with dirty nails
<p><b>Scabies</b></p> <p>Image Source: University of Washington Seattle STD/HIV Prevention Training Centre</p>		"The Itch"	10 months	Itching at night. Red lines in the skin as the scabies burrow. Ulcers develop after scratching.	Special cream and soap. All clothing and bedding must be washed in hot, soapy water.	Spreads all over the body, secondary infection of the skin due to scratching with dirty nails

**MEN'S STI's (PAGE 2)**

STI	IMAGE	KNOWN AS	COURSE	SYMPTOMS	TREATMENT	WITHOUT TREATMENT
<p><b>Genital Herpes</b></p> <p>Image Source: University of Washington Seattle STD/HIV Prevention Training Centre</p>		<p>Blisters</p>	<p>2-20 days before presenting with symptoms</p>	<p>Type 1-fever blisters on lips diseases above waist. Type 2-diseases below the waist genital herpes. Genital herpes can be purely genital or oral genital due to oral sex. Fever, swollen glands, frequent urination. Male lesions are on the prepuce or shaft of penis, small fluid-filled blisters in the genital and anal area</p>	<p>Anti-viral treatment in severe cases, remember drugs only alleviate the symptoms it doesn't cure the problem, will be recurrent.</p>	<p>Lesions/blisters might disappear but re-occur at a later stage</p>
<p><b>Non-specific urethritis</b></p> <p>Image Source: SOA-AIDS Amsterdam</p>		<p>NSU</p>	<p>1-7 days</p>	<p>Men: Burning urination and discharge</p>	<p>Antibiotics</p>	<p>Severe infection of reproductive organs and infertility</p>
<p><b>Genital warts</b></p> <p>Image Source: University of Washington Seattle STD/HIV Prevention Training Centre</p>		<p>Warts</p>	<p>1-6 months</p>	<p>Genital warts develop in moist areas and can become malignant. Small painless bumps growing on genitals with itching and burning. Maybe inside the urethra of men</p>	<p>Removed by burning, freezing or minor surgery. Special lotion</p>	<p>Lead to cervical cancer. Grow large and spread.</p>
<p><b>Hepatitis B</b></p>		<p>Hep B – Blood transfusion, sexually transmitted, saliva &amp; needles</p>	<p>Symptoms develop within 30-180 days of exposure to the virus. The symptoms are often compared to flu.</p>	<p>Appetite loss, Feeling tired, nausea and vomiting, itching all over the body, pain over the liver (on the right side of abdomen under the rib cage), yellowish skin tone and eyes, dark urine (like tea) and pale grey stools</p>	<p>If very severe, symptoms such as vomiting or diarrhoea may require treatment to restore fluids and electrolytes. No cure. To prevent, Hep B vaccine</p>	<p>Can lead to permanent liver damage and even cancer.</p>
<p><b>Chancroid</b></p> <p>Image Source: CDC/Joe Miller 1974</p>			<p>1-7 days</p>	<p>First sign are one or more sores filled with puss and surrounded by red border, eventually ruptures leaving painful open wounds, swollen glands</p>	<p>Antibiotics</p>	<p>Severe infection in the genital area, can even lead to amputation of penis</p>
<p><b>Gonococcal Pharyngitis</b></p> <p>Image Source: Public Health Image Library for Centre for Disease Control and Prevention: Drs N.J. Flumara and Gavin Hart</p>			<p>1-7 days</p>	<p>Difficulty in swallowing, fever, sore throat, swollen glands in the neck, abnormal sense of taste, headache, nasal congestion, nasal discharge, muscle and joint pain and even skin lesions</p>	<p>Antibiotics</p>	<p>Severe infection of the throat and the disease can spread to other regions of the body</p>