











WOMEN'S STI's (PAGE 1)

STI	IMAGE	KNOWN AS	COURSE	SYMPTOMS	TREATMENT	WITHOUT TREATMENT
Syphilis Image Source: University of Washington Seattle STD/HIV Prevention Training Centre		The Pox	Stage 1: 1-3 months Stage 2: 3-6 months Stage 3: Many years	Stage 1: A painless sore called a chancre. Stage 2: Fever, headache and general rash. Stage 3: Very ill. The cause is not easy to find.	Antibiotics	Severe infection, infertility, skin diseases, arthritis, baby can be born blind.
Gonorrhoea		Drip, Clap, Dose	1 – 10 days	Most women have pelvic pain, painful urination, vaginal discharge or fever.	Antibiotics	Vaginal infection
Trichomoniasis Image Source: University of Washington Seattle STD/HIV Prevention Training Centre		Trich	1 week	Yellow or greenish frothy discharge, itching around the genitals.	Antibiotics	Fever, infection of the organs, infection passed on to baby
Chlamydia Image Source: University of Southern Mississippi Health.Promotion@usm.edu		PID	1-3 weeks	Pelvic pain, vaginal discharge, painful and frequent urination, bleeding after sexual intercourse or no symptoms.	Antibiotics	Severe infection of reproductive organs.
Non-Gonococcal urethritis		NGU	1-3 weeks	Frequent painful urination and a discharge.	Antibiotics	Infertility
Candida (yeast) Image Source: University of Washington Seattle STD/HIV Prevention Training Centre		Thrush, yeast	1-7 days	Thick white discharge, swelling of vulva, painful and frequent irritation, itching around the genitals.	Vaginal cream	Extreme discomfort
Pubic Lice Image Source: University of Washington Seattle STD/HIV Prevention Training Centre		Crabs	Immediately	Itching in area of the chest, genital hair. Lice crawling and small eggs on hair or clothing.	Special shampoos or lotions. All clothing and bedding must be washed in hot, soapy water.	Irritation
Scabies		"The Itch"	10 months	Itching at night. Red lines in the skin as the scabies burrow. Ulcers develop after scratching.	Special cream and soap. All clothing and bedding must be washed in hot, soapy water.	Spreads all over the body, secondary infections on the skin due to scratching with dirty nails

WOMEN'S STI's (PAGE 2)

STI	IMAGE	KNOWN AS	COURSE	SYMPTOMS	TREATMENT	WITHOUT TREATMENT
<p>Genital herpes</p> <p>Image Source: SOA-AIDS Amsterdam</p>		<p>Blisters</p>	<p>2-20 days before presenting with symptoms</p>	<p>Type 1-fever blisters on lips diseases above waist. Type 2-diseases below the waist genital herpes. Genital herpes can be purely genital or oral genital due to oral sex. Fever, swollen glands, frequent urination vaginal discharge or bleeding. Female's lesion labia majora or minora, buttocks or thighs.</p>	<p>Anti-viral treatment is given in severe cases, remember drugs only alleviate the symptoms it doesn't cure the problem, can be recurrent.</p>	<p>Lesions/blisters might disappear but recur at a later stage</p>
<p>Pelvic inflammatory Disease</p> <p>Image Source: PDRhealth.com</p>		<p>PID</p>		<p>Fever, nausea, vomiting, lower abdominal pain, painful sex and painful menstruation</p>	<p>Antibiotics</p>	<p>Sterility and abscess of the tubes</p>
<p>Non-specific urethritis</p> <p>Image Source: SOA-AIDS Amsterdam</p>		<p>NSU</p>	<p>1-7 days</p>	<p>Women: Fewer symptoms, burning urine and frequent urination</p>	<p>Antibiotics</p>	<p>Severe infection of reproductive organs and infertility.</p>
<p>Genital warts</p> <p>Image Source: Brooklyn Gyn Place – Prof Erwin J. Haeberle</p>		<p>Warts</p>	<p>1-6 months</p>	<p>Genital warts develop in moist areas and can become malignant. Small painless bumps growing on genitals with itching and burning. May be inside the vagina and on cervix Need to do a pap smear regular</p>	<p>Removed by burning, freezing or minor surgery. Special lotion</p>	<p>Lead to cervical cancer. Grow large and spread.</p>
<p>Hepatitis B</p>		<p>Hep B – Blood transfusion, sexually transmitted, saliva & needles</p>	<p>Symptoms develop within 30-180 days of exposure to the virus. The symptoms are often compared to flu.</p>	<p>Appetite loss, Feeling tired, nausea and vomiting, itching all over the body, pain over the liver (on the right side of abdomen under the rib cage), yellowish skin tone and eyes, dark urine (like tea) and pale grey stools</p>	<p>If very severe, symptoms such as vomiting or diarrhoea may require treatment to restore fluids and electrolytes. No cure. To prevent, Hep B vaccine</p>	<p>Can cause permanent liver damage and liver cancer</p>